

B'  
binding groups include, but are not limited to, boronic acid, crown ether, and aza-crown ether, such as 1,4,7,10,13-Pentaoxa-16-aza-cyclooctadecane (aza 18-crown-6) and 1,4,7,13-tetraoxa-10-aza-cyclohexadecane (aza 15-crown-5). In a preferred embodiment, the Bd1 is R1-B(OH)<sub>2</sub> and Bd2 is R2-B(OH)<sub>2</sub>. R1 and R2 are aliphatic or aromatic functional groups selected independently from each other and B is a boron atom.

Please replace the text of the first full paragraph on page 8 with the following text:

Ba  
202704225904  
In the present invention, the binding groups may be any functional groups, as long as they provide the desired specific binding of the analyte to the sensor with a formation of 1:1 complex. The binding groups are preferably electron deficient groups. The electron deficiency governs the shift of the unshared electron pair from the nitrogen atoms to the binding group when specifically binding the analyte. Examples of the acceptable binding groups include, but are not limited to, boronic acid, crown ether, and aza-crown ether, such as 1,4,7,10,13-Pentaoxa-16-aza-cyclooctadecane (aza 18-crown-6) and 1,4,7,13-tetraoxa-10-aza-cyclohexadecane (aza 15-crown-5). Examples of analytes that may be identified by utilizing sensors of the present invention include, but are not limited to, saccharides, amino saccharides, and carbonyl saccharides.

IN THE CLAIMS:

Please cancel claims 1-21 without prejudice.

Please replace the text of claim 33 with the following text: